

Post-Operative Instructions

Shoulder Arthroscopy

Activity

- Wear your sling at all times until the nerve block has completely resolved. Be sure to support your wrist in the sling. Based upon your surgery:
 - You must remain in your sling at all times except when showering. The only exception to this is to allow you to stretch your elbow a few times a day to prevent your hand and forearm from swelling. Do NOT use your arm to reach for, lift or carry anything until instructed otherwise.
 - You may remove your sling when the arm is comfortable. You may use the arm as tolerated.
- Formal physical therapy will begin after your first surgical follow-up visit.

Pain Control

- You have been prescribed a pain medication that contains a narcotic. Do not drink alcohol, drive a car or operate heavy machinery while on this medication.
 - Do NOT take Tylenol (acetaminophen) while on this medication as it may cause an overdose.
- You may take anti-inflammatory medications such as Ibuprofen or Naprosyn (with food) in addition to the pain medication.
- Apply ice to the shoulder for 15 minutes as often as every hour. You may continue this for as many days as you like.
- You may find sleeping on an incline (reclining chair) to be more comfortable for the first few days.

Wound Care

- **You may remove the bandages and shower in 48 hours.**
 - Do NOT bathe or go swimming until the sutures are removed.
- Cover the incisions with band-aids after your shower.
- Do not apply any creams or lotions to the incisions.

Follow-up Appointment

- Your follow-up appointment should be 10-14 days after surgery, please call the office if you do not have an appointment. We will review the findings from your surgery and remove your sutures during this visit.
- Please call the office with any questions or concerns, or if you experience any of the following: fevers (greater than 101°F), chills, increased swelling, pain, redness or increasing drainage at the incision.