

Post-Operative Instructions

Trigger Finger Release

Activity

- Leave your dressing on at all times. You can use a sling for comfort.
 - Do NOT use your hand to lift or carry anything (ie. **non-weight bearing**) until instructed otherwise.
- **Elevate your operative hand on pillows for the first 72 hours.**
 - To reduce swelling, your hand must be elevated above the level of your heart.
 - You may feel increased pain or pressure any time the hand drops below the level of your heart.
- Formal physical therapy will begin after your first surgical follow-up visit.

Pain Control

- You have been prescribed a pain medication that contains a narcotic. Do not drink alcohol, drive a car or operate heavy machinery while on this medication.
 - Do NOT take Tylenol (acetaminophen) while on this medication as it may cause an overdose.
- You may take anti-inflammatory medications such as Ibuprofen or Naprosyn (with food) in addition to the pain medication.
- Apply ice to the hand for 15 minutes as often as every hour. You may continue this for as many days as you like.

Wound Care

- **Do not remove your surgical dressing. Keep it clean and dry.** It will be changed during your postoperative evaluation.

Follow-up Appointment

- Your follow-up appointment should be 10-14 days after surgery, please call the office if you do not have an appointment. We will review the findings from your surgery and remove your sutures during this visit.
- Please call the office with any questions or concerns, or if you experience any of the following: fevers (greater than 101°F), chills, increased swelling, pain, redness or increasing drainage at the incision.